

## Case Report

## Improved Insulin Resistance and Glucose Variability by Super-Low Carbohydrate Diet

Hirohisa Urasaki, BSc<sup>1,2</sup>; Hiroshi Bando, MD, PhD, FACP<sup>2,3,4\*</sup>; Harumi Kobayashi, RN<sup>4</sup>; Hiroko Urasaki, BSc<sup>2</sup>; Hiroko Ogawa, PharmD<sup>4</sup>; Miwako Nakanishi, MD, PhD<sup>4</sup>; Osami Watanabe, MD, PhD<sup>4</sup>; Shinji Nagahiro, MD, PhD<sup>4,5</sup>

<sup>1</sup>Nankai Hospital, Naruto City, Tokushima, Japan

<sup>2</sup>Tokushima Low Carbohydrate Diet Promotion Association, Tokushima, Japan

<sup>3</sup>Instructor, Tokushima University, Tokushima, Japan

<sup>4</sup>Yoshinogawa Hospital, Tokushima, Japan

<sup>5</sup>Emeritus Professor, Former Director of Tokushima University Hospital, Tokushima, Japan

## \*Corresponding author

Hiroshi Bando, MD, PhD, FACP

Instructor, Tokushima University, Nakashowa 1-61, Tokushima 770-0943 Japan; Tel. +81-90-3187-2485; E-mail: [pianomed@bronze.ocn.ne.jp](mailto:pianomed@bronze.ocn.ne.jp)

## Article information

Received: February 25<sup>th</sup>, 2022; Revised: March 5<sup>th</sup>, 2022; Accepted: March 7<sup>th</sup>, 2022; Published: March 7<sup>th</sup>, 2022

## Cite this article

Urasaki H, Bando H, Kobayashi H, et al. Improved insulin resistance and glucose variability by super-low carbohydrate diet. *Diabetes Res Open J.* 2022; 8(1): 11-17. doi: [10.17140/DROJ-8-154](https://doi.org/10.17140/DROJ-8-154)

## ABSTRACT

## Background

Diabetes mellitus (DM) has been more prevalent. American Diabetes Association (ADA) proposed the Standards of Medical Care in Diabetes-2022. For nutritional therapy, low carbohydrate diet (LCD) has been recognized for its benefits. Authors have continued diabetic research concerning LCD and meal tolerance test (MTT).

## Case Presentation

The case is 61-year-old male with type 2 diabetes mellitus (T2DM) for years. His hemoglobin A1c (HbA1c) increased to 7.8% in autumn 2021, and further evaluation and treatment was conducted including LCD, daily check of meal and carbohydrate amount, 75 g OGTT, glucagon stimulation test (GST) and others.

## Results

He was on super-LCD method including 12% of carbohydrate. His carbohydrate intake amount and 45-minutes post-prandial blood glucose showed significant correlation. The results of 75 g OGTT twice in May 2020 and December 2021 showed that similar pattern of glucose and insulin responses and insulinogenic index (IGI). In contrast, they showed decreased fasting immuno-reactive insulin (IRI) and Homeostasis model assessment insulin resistance (HOMA-R). For GST, C-peptide showed normal response.

## Discussion and Conclusion

Judging from the results of MTT, OGTT, GST and IGI, he seems to show rather decreased insulin resistance by LCD associated with preserved insulin secretion ability to some degree. Further investigation would be required from pathophysiological point of view.

## Keywords

Super-low carbohydrate diet (LCD); Meal tolerance test (MTT); Type 2 diabetes mellitus (T2DM); Insulinogenic index (IGI); Glucagon stimulation test (GST); Homeostasis model assessment insulin resistance (HOMA-R).

## INTRODUCTION

Diabetes mellitus (DM) has been more prevalent across the world according to the epidemiological report of International Diabetes Federation (IDF).<sup>1</sup> Among them, undiagnosed DM (UDM) in adult is gradually increasing in developed and developing countries.<sup>2</sup> American Diabetes Association (ADA) proposed the latest guideline in Jan 2022, which is the Standards of Medical Care in Diabetes-2022.<sup>3</sup>

For the diabetic therapy, nutritional treatment would be the fundamental strategy including low carbohydrate diet (LCD).<sup>4</sup> Clinical efficacy of LCD has been gradually recognized for improving glucose variability and weight reduction.<sup>5</sup> Authors and collaborators have continued diabetic research and practice for long.<sup>6</sup> Their clinical fields include LCD, continuous glucose monitoring (CGM), Glucagon-Like Peptide 1 receptor agonist (GLP-1RA), meal tolerance test (MTT) and others.<sup>7,8</sup>

Concerning LCD investigation, historical changes for nu-

tritional therapy were observed. Calorie restriction (CR) was the standard method for diabetic diet therapy in previous period. After that, LCD method was introduced to medical and health care field by Atkins et al.<sup>9</sup> LCD was evaluated to be clinically effective by Dietary Intervention Randomized Controlled Trial (DIRECT) study.<sup>10</sup> Successively, LCD was generally defined as related with several diet methods.<sup>11</sup> On the other hand, authors et al. initiated LCD in Japan and developed LCD medically and socially through the activity of Japan LCD Promotion Association (JLCDPA).<sup>12</sup> Among our various activities, three representative methods of LCD were widely informed to general people. They are petite LCD, standard LCD and super LCD.<sup>13</sup> These LCD nutritional options were applied to many obesity patients, and the results showed that 25% subjects achieved >10% of weight reduction.<sup>14</sup>

As to MTT, standardized liquid meal was ingested and assessed for  $\beta$ -cell function in diabetic patients.<sup>15</sup> Similar to MTT, widely applied popular method includes oral glucose tolerance test (OGTT).<sup>16</sup> Authors have developed MTT research, and proposed the simple method using breakfast.<sup>17</sup> The standard meal is made from Japanese style standard breakfast including 70 g of carbohydrate.<sup>18</sup> Applying this meal, the responses of blood glucose, insulin and C-peptide were investigated.<sup>19</sup>

By combination of the research of LCD and MTT, we have continued clinical diabetic studies. Among several patients with type 2 DM (T2DM), we have experienced a case of impressive characteristic aspects. In this case report, general outline of clinical course and treatment will be introduced.

## CASE PRESENTATION

### Family and Past History

The current case is 61-year-old man. Regarding his family history, there was unremarkable disease involved, such as diabetes mellitus, hypertension, coronary heart disease (CHD), cerebral vascular accident (CVA) or malignant disease.

For his previous medical history, he was born without any health problem. When he was born, his weight was 3500 g. It was much heavier than average baby weight. Her mother was not diabetic or did not have any medical problems during pregnancy. He did not have significant disease or impaired function until 54-years-old.

### Medical History and Status

The current case is 61-year-old T2DM male patient who has diabetic history of 4-years. He has become gradually rather obese after 55 years old and then he was diagnosed for T2DM. Then he was advised to reduce the weight, and he tried to continue mild LCD for a few years. He received 75 g OGTT in May 2020, in which it showed delayed pattern of glucose and insulin. The HbA1c value maintained about 6.1-6.4% for a few years. During summer 2021, he felt general malaise and weight loss and then received medical check. Consequently, he was pointed out to have exacerbated T2DM and to start strict LCD at once. Consequently, he and his wife have started to evaluated carbohydrate amount of his daily

meal. Consequently, HbA1c and blood glucose decreased, associated with lower post-prandial blood glucose.

His physical status showed unremarkable findings, including consciousness, vital signs, lung, heart, abdomen and neurological examination. His stature, weight and body mass index (BMI) was 181 cm, 89 kg and 27.2 kg/m<sup>2</sup>, respectively. His laboratory examinations revealed that HDL-C 65 mg/dL, LDL-C 142 mg/dL, TG 100 mg/dL, GOT 22 U/L, GPT 28 U/L,  $\gamma$ -GTP 25 U/L, BUN 17 mg/dL, Cr 0.8 mg/dL, UA 3.9 mg/dL, WBC 8600 / $\mu$ L, RBC 508 $\times$ 10<sup>6</sup> / $\mu$ L, Hb 16.2 g/dL, Plt 21.1 $\times$ 10<sup>4</sup> / $\mu$ L, CRP 0.02 mg/dL. Urinalysis showed pH 6.0, protein (-), glucose (+), urobilinogen (+/-), occult blood (-) and ketone body (+). Other tests showed negative for chest X-P and electrocardiogram (ECG). Pulse wave velocity (PWV) test showed normal range results, where ankle brachial index (ABI) was 1.08/1.09 and cardio-ankle vascular index (CAVI) 10.0/10.0 for right/left.

### Clinical Course

This case has applied strict LCD that has been known as super-LCD method. It means that carbohydrate ratio in calorie calculation would be 12%. He actually continued to decrease carbohydrate as possible.<sup>20</sup> Then, his daily profile of blood glucose became normalized and HbA1c values decreased from 7.8% to 6.3% for 3-months. He felt no remarkable complaints, symptoms, signs or abnormal biomedical exams during 4-months. During his clinical course, he received 75 g OGTT and continued taking daily meal data by digital camera associated with calculating carbohydrate amount and calorie included. Especially, 45-minutes post-prandial glucose of dinner was continued to check for months.<sup>21</sup> Furthermore, the insulin response was investigated for endocrinological stimulus by glucagon test.

## RESULTS


During the period of LCD, several data such as carbohydrate amount and calorie were checked and analyzed (Table 1). Table 1 revealed Western and Japanese styles of dinner provided. Blood glucose value was measured as 45-minutes post-prandial point of each dinner. Further, relationship between carbohydrate amount and 45-minutes post-prandial blood glucose was investigated, which was analyzed in Figure 1. It shows significant relationship between them. The formula for the regression curve is shown, which was  $y=1.9084x+104.76$ ,  $R^2=0.5789$ ,  $p<0.05$ .

Regarding the ability of insulin secretion, 75 g OGTT was conducted (Figure 2). Figure 2 includes totally four results of 75 g OGTT, in which Figures 2a and 2b are from the first author of current T2DM case in May 2020 and December 2021 (case 1), Figure 2c is from the second author without any diseases (case 2) and Figure 2d is from the third author with T2DM (case 3). Case 1 showed delayed response of glucose and immuno-reactive insulin (IRI) for T2DM. Case 2 showed normal response of glucose and IRI for normal subject. Case 3 showed hyperglycemia in which glucose increased to 240 mg/dL in 120-minutes. Related to 4 data of Figure 2, detail biomarkers of diabetes mellitus were summarized associated with detail background of case 1-3 in Table 2. Case 1 showed elevated fasting IRI, homeostasis model assessment insu-

**Table 1: Dinner Content with Carbohydrate Amount and Post-Prandial Blood Glucose**

Dinner	Ingredient	Each Carbo (gram)	Total Carbo (gram)	Total Calorie (kcal)
<b>A</b> 	Tempura	21.2	48.2	656
	Fish	4.0		
	White rice	19.0		
	Miso soup, etc.	4.0		
<b>B</b> 	Fried chicken	28.0	33.4	652
	Cauliflower	0.1		
	Vegetables	0.1		
	Miso soup, etc.	5.4		
<b>C</b> 	Hot pot shabu-shabu	0.1	10.8	445
	Beef	0.3		
	Tofu	0.4		
	Vegetables	10.0		
<b>D</b> 	Chili pepper soup	7.9	30.0	205
	Pork	0.2		
	White radish	2.4		
	Vegetables	19.5		



**Figure 1. Relationship between Carbohydrate Amount in the Dinner and 45-minutes Post-prandial Blood Glucose**



**Table 2: General Background, HOMA Values and IGI in Three Cases**

Case	Age/ Sex	Diagnosis	Test date (Mo/Year)	FPG (mg/dL)	IRI (μIU/L)	HOMA-R	HOMA-1β (%)	IGI	Figure
Case 1	61 M	T2DM	May 2020	132	11.6	3.78	60.5	0.04	Figure 2a
Case 1	61 M	T2DM	Dec 2021	115	5.5	1.56	38.1	0.03	Figure 2b
Case 2	65 M	Normal	Jan 2022	83	2.7	0.55	48.6	1.76	Figure 2c
Case 3	56 F	T2DM	Feb 2022	119	3.7	1.09	23.8	0.12	Figure 2d

Case 1 is the first author who works in the administration section of the hospital.  
Case 2 is the second author who specializes diabetes mellitus as a physician.  
Case 3 is the third author who has worked in the out clinic as a hospital nurse.



lin resistance (HOMA-R) in May 2020 and almost normal range of IRI and HOMA-R in December 2021. His insulinogenic index (IGI) values remained low during the period. Case 2 is 65-year-old physician and showed normal range of all biomarkers. Case 3 is 56-year-old nurse and showed normal range of HOMA-R and Homeostatic model assessment of  $\beta$ -cell function (HOMA- $\beta$ ), and decreased IGI. For detection of insulin secretion, glucagon stimulation test (GST) was conducted. As a result, c-peptide value increased from 2.0 ng/mL to 4.7 ng/mL during 6-minutes (Figure 3).

## DISCUSSION AND CONCLUSION

The current case has T2DM, and HbA1c value has been elevated for years. Regarding the pathophysiological aspect, both of insulin resistance and insulin deficiency has been present to some degree from his history and biomedical examination results. BMI is currently 27.2 kg/m<sup>2</sup>, but once it weighed 100 kg with 30.5 kg/m<sup>2</sup> of BMI. In this case report, detail investigation for several aspects is conducted, which are summarized in tables and figures. Discussion and perspectives would be described in this order.

In this case, super low carbohydrate diet (super-LCD) was applied and continued.<sup>13</sup> It includes 12% of carbohydrate in caloric

ratio, associated with normalized HbA1c and glycemic fluctuation. Along with super-LCD, he successfully continued meal diary every day and blood glucose profile perfectly (Table 1). We have continued to research and clinically apply LCDs for long. Among them, the clinical effect of LCD was found in many patients, and the cases of remarkable weight loss were also observed.<sup>20</sup> There are actually three methods for LCDs, which are petite-LCD, standard-LCD and super-LCD. The carbohydrate content in caloric ratio is 12%, 26% and 40%, respectively.<sup>13</sup> From recent reports on LCD, various beneficial effects have been observed.<sup>22</sup>

For evaluating clinical LCD trials, carbohydrate ratio in the meal has been on discussion.<sup>17</sup> A standard classification shows that high-carbohydrate diet (>45%), moderate carbohydrate diet (26-45%), LCD (10-25%), and very low-carbohydrate ketogenic diet (VLCKD) (<10%).<sup>18,19</sup> As carbohydrate ratio is lower, glucose-lowering efficacy is larger for T2DM.<sup>20</sup> However, it has been not clear whether more carbohydrate restriction can bring the cardiovascular risk reduction or not.<sup>19</sup>

Meal tolerance test (MTT) was performed in advance to examine post-prandial blood glucose 30-minutes, 45-minutes, and 60-minutes for the case. As a result, 45-minutes post-prandial glucose was the highest.<sup>21</sup> Then, post-prandial timing was de-

cided 45-minutes for this case. Significant correlation was found between carbohydrate amount and 45-minutes post-prandial glucose in the dinner (Figure 1). We can calculate using the regression curve. When 20 g of carbohydrate is input, glucose value becomes 142 mg/dL, and when 50 g is input, it becomes 199 mg/dL. The measurement seems to be more or less consistent with actual carbohydrate intake and post-prandial glucose in T2DM patients.<sup>18</sup> MTT and OGTT are often compared for simple and convenient in the clinical practice.<sup>23</sup> Both are suitable methods to evaluate physiological  $\beta$ -cell ability for long-term.<sup>24</sup> Comparative investigation revealed that MTT showed stronger response of  $\beta$ -cell than OGTT,<sup>24</sup> and represented reproducible and reliable way of analyzing biomarkers of  $\beta$ -cell function.<sup>25</sup>

OGTT exams were checked twice, which were compared between in May 2020 and December 2021 (Figure 2). His weight was 95 kg *vs* 88 kg and his fasting IRI was 11.6  $\mu$ IU/mL *vs* 5.5  $\mu$ IU/mL. From these data, the degree of insulin resistance may be reduced. The pattern of glycemic and IRI for 75 g OGTT seems to be similar, and unremarkable change in insulin secretory reserve was observed. As a comparative control, case 2 showed rather low glucose on 120-minutes. Case 2 is healthy physician and usually on standard LCD, which may be involved in lower glucose on 120-min after 75 g OGTT. Case 3 has T2DM for more than 15-years, and is currently on petite LCD for better glucose variability. It showed delayed insulin response and blood glucose increased from 119 mg/dL to 334 mg/dL (215 mg/dL increase). From pathophysiological point of view, T2DM patient is reported to show 3 mg/dL of glucose increase per 1 g intake of carbohydrate. When calculating as 75 g $\times$ 3=225, this result seems to be similar to actual data.<sup>21</sup> OGTT has been widely used for investigating glucose tolerance and insulin secretion in clinical practice.<sup>26</sup> Estimation of  $\beta$  cell function by OGTT shows more physiological responses of glucose and insulin than those of frequently sampled intravenous glucose tolerance tests (FSIVGTT) and hyperglycemic clamp.<sup>27</sup>

HOMA-R, HOMA- $\beta$  and IGI in case 1 suggested that insulin resistance was slightly reduced for 19-months, but insulin secretory capacity was similar (Table 2). Regarding the pathogenesis of all types of DM, increased insulin resistance and reduced insulin secretion would play a major role.<sup>26</sup> It is rather difficult to obtain complete pathophysiology of diabetic metabolism because glucose and insulin have been influenced through various complex process. However, some metabolic indices are appropriate for investigate clinical diabetic situation, which are HOMA-R, HOMA- $\beta$  and IGI. Regarding these biomarkers, the standard values widely used so far are HOMA-R<2.5, HOMA- $\beta$ >30%, and IGI>0.4-0.5.

The standard range of HOMA-R cannot be clearly defined, where 2.5 has been often used in usual situation. The cut-off point of HOMA-R was studied in reference to the definition of IDF and Adult Treatment Panel III (ATPIII) criteria.<sup>28</sup> The method included the analysis of area under the curve (AUC). As a result, the optimal cut-off of HOMA-R was 1.775 for diagnosis of Metabolic syndrome (MetS) in non-DM patients, 4.325 and 3.875 for diagnosis of DM in MetS patients for IDF and ATPIII, respectively. Consequently, MetS or diabetes can be diagnosed from various patients with or without diabetes in applying this optimal

method. HOMA- $\beta$  has been one of the useful static assessments of  $\beta$ -cell function.<sup>29</sup> It is simple and widely applied in epidemiologic investigation. However, it has some limitation because only one point of glucose and insulin cannot represent complex situation of glucose-insulin feedback axis.<sup>30</sup>

Regarding IGI, lower results were found in case 1 and 4, and normal in case 2 (Table 2). Case 1 has weight reduction compared to 19 months ago, leading to decreased fasting IRI and HOMA-R. However, the capacity of insulin secretion did not change. IGI has been widely used to quantify the response of  $\beta$ -cell for the changes in blood glucose. It is calculated for delta-IRI (0-30min) ( $\mu$ IU/mL)/delta-glucose (0-30-minutes) (mg/dL).<sup>24</sup> It is also applied in the case of MTT trials.<sup>31</sup> Compared IGI for GTT and MTT has been studied.<sup>32</sup> Authors have studied insulin and C-peptide immunoreactivity (CPR) responses for MTT and presented clinical efficacy of MTT.<sup>19,21</sup> From several reports,  $\beta$ -cell function seems to be higher during MTT than OGTT, suggesting that several function parameters would be influenced by some composition in the meal.<sup>24</sup>

Glucagon stimulation test (GST) has been applied for the investigation of insulin-secreting ability from  $\beta$ -cell in the pancreas. Formerly, the purpose was evaluation of residual  $\beta$ -cell function,<sup>33</sup> in which subjects were 17 insulin-dependent diabetes mellitus (IDDM) patients. C-peptide elevation was measured after meal ( $p<0.01$ ) and after intravenous glucagon ( $p<0.001$ ). Both responses showed correlation ( $p<0.005$ ), suggesting possible prediction of glucagon test for the responsive ability of  $\beta$ -cells during usual daily life. In the current case, GST revealed that pancreatic  $\beta$ -cell function was preserved (Figure 3). Based on his data for clinical course, 75 g OGTT, HOMA-R, HOMA- $\beta$ , IGI and glucagon tests were comprehensively analyzed totally. LCD continuation had successfully brought weight reduction, decreased HbA1c, moderately reduced insulin resistance. Furthermore, it seems to maintain secretory capacity against strong endocrinological stimulation. On the other hand, OGTT seems to be rather weaker stimulation compared to GST. IGI values by 75 g OGTT did not improve.

There are some limitations for this report as follows: i) super LCD was continued, but any food has carbohydrate to some degree for elevating blood glucose, ii) correlation between carbohydrate amount and post-prandial glucose was found, but the study of MMT cannot predict complete suggested glucose values due to ingestion the meal per OS, iii) OGTT twice showed delayed IRI responses, which may suppose persisting impaired insulin secretion and increased insulin resistance, iv) current glucose-insulin situation will be possible changed due to clinical course in the future, then detail follow-up the glucose variability would be required.

In summary, this case report described a male patient with T2DM and gave some discussion from several points of view. He successfully continued daily check of meal with carbohydrate amount and post-prandial blood glucose. Diabetic biomarkers including 75 g OGTT, HOMA-R, HOMA- $\beta$ , IGI and GST were analyzed for evaluating current pathophysiology. Current information will become hopefully some reference for diabetic practice and research in the future.

## ETHICAL CONSIDERATIONS

Current investigation was conducted along the Declaration of Helsinki, that was previously revised in 2013 for the WMA Fortaleza General Assembly. In addition, several commentaries were added by the ethical guidelines for medical research. They are notified by the Ministry of Education, Culture, Sports, Science and Technology (MEXT), Japan and Ministry of Health, Labour and Welfare (MHLW), Japan. This study in detail was explained to the patient. Authors have obtained the written document agreements from the patient. Current study was discussed in the professional ethical committee. The committee involves several professionals including president, director, doctors, nurses, pharmacists, dieticians, and a professional legal specialty.

## FUNDING

There was no funding received for this paper.

## CONFLICTS OF INTEREST

The authors declare that they have no conflicts of interest.

## REFERENCES

- Aschner P, Karuranga S, James S, et al. The international diabetes federation's guide for diabetes epidemiological studies. *Diabetes Res Clin Pract.* 2021; 172: 108630. doi: [10.1016/j.diabres.2020.108630](https://doi.org/10.1016/j.diabres.2020.108630)
- Ogurtsova K, Guariguata L, Barengo NC, et al. IDF diabetes Atlas: Global estimates of undiagnosed diabetes in adults for 2021. *Diabetes Res Clin Pract.* 2022; 183: 109118. doi: [10.1016/j.diabres.2021.109118](https://doi.org/10.1016/j.diabres.2021.109118)
- American Diabetes Association. Introduction: Standards of medical care in diabetes—2022. *Diabetes Care.* 2022; 45(Suppl. 1): S1-S2. doi: [10.2337/dc22-S1NT](https://doi.org/10.2337/dc22-S1NT)
- Gram-Kampmann EM, Hansen CD, Hugger MB, et al. Effects of a 6-month, low-carbohydrate diet on glycaemic control, body composition, and cardiovascular risk factors in patients with type 2 diabetes: An open-label randomized controlled trial. *Diabetes Obes Metab.* 2022. doi: [10.1111/dom.14633](https://doi.org/10.1111/dom.14633)
- ADA Professional Practice Committee. 5. Facilitating behavior change and well-being to improve health outcomes: Standards of Medical Care in Diabetes-2022. *Diabetes Care.* 2022; 45 (Suppl. 1): S60-S82. doi: [10.2337/dc22-S005](https://doi.org/10.2337/dc22-S005)
- Bando H, Iwatsuki N, Ogawa T, Sakamoto K. Investigation for daily profile of blood glucose by the administration of canagliflozin and xultophy (Ideglira). *Int J Endocrinol Diabetes.* 2022; 5(1): 129. doi: [10.36266/IJED/129](https://doi.org/10.36266/IJED/129)
- Iwatsuki N, Bando H, Okada M. Pharmacological characteristic of imeglimin (Twymeeg) for dual mechanism to insulin secretion and resistance. *SunText Rev Pharm Sci.* 2022; 3(1): 113. doi: [10.51737/2766-5232.2022.013](https://doi.org/10.51737/2766-5232.2022.013)
- Hayashi K. Useful xultophy for older diabetic with various problems. *SunText Rev Med Clin Res.* 2021; 2(2): 126. doi: [10.51737/2766-4813.2021.027](https://doi.org/10.51737/2766-4813.2021.027)
- Atkins RC. *Dr. Atkins' New Diet Revolution: The No-hunger, Luxurious Weight Loss Plan That Really Works!* Harper. London, UK: Ebury Digital; 2009.
- Shai I, Schwarzfuchs D, Henkin Y, et al. Weight loss with a low-carbohydrate, Mediterranean, or low-fat diet. *N Engl J Med.* 2008; 359(3): 229-241. doi: [10.1056/NEJMoa0708681](https://doi.org/10.1056/NEJMoa0708681)
- Feinman RD, Pogozelski WK, Astrup A, et al. Dietary carbohydrate restriction as the first approach in diabetes management: critical review and evidence base. *Nutrition.* 2015; 31(1): 1-13. doi: [10.1016/j.nut.2014.06.011](https://doi.org/10.1016/j.nut.2014.06.011)
- Ebe K, Ebe Y, Yokota S, et al. Low carbohydrate diet (LCD) treated for three cases as diabetic diet therapy. *Kyoto Medical Association Journal.* 2004; 51: 125-129.
- Bando H. Useful tips for actual low carbohydrate diet (LCD) with super-, standard- and petite-LCD methods. *EC Nutrition.* 2020; 15(5): 01-04.
- Nakamura T, Kawashima T, Dobashi M, Narita A, Bando H. Effective nutritional guidance for obesity by low carbohydrate diet (LCD). *Asp Biomed Clin Case Rep.* 2019; 2(s1): 16-21. doi: [10.36502/2019/ASJBCCR.6141](https://doi.org/10.36502/2019/ASJBCCR.6141)
- Besser REJ, Shields BM, Casas R, Hattersley AT, Ludvigsson J. Lessons from the mixed-meal tolerance test: Use of 90-minute and fasting C-peptide in pediatric diabetes. *Diabetes Care.* 2013; 36: 195-201. doi: [10.2337/dc12-0836](https://doi.org/10.2337/dc12-0836)
- Kössler T, Bobrov P, Strassburger K, et al. Impact of mixed meal tolerance test composition on measures of beta-cell function in type 2 diabetes. *Nutr Metab (Lond).* 2021; 18: 47. doi: [10.1186/s12986-021-00556-1](https://doi.org/10.1186/s12986-021-00556-1)
- Bando H, Ebe K, Muneta T, Bando M, Yonei Y. Proposal for insulinogenic index (IGI)-Carbo70 as experimental evaluation for diabetes. *J Clin Exp Endocrinol.* 2017; 1: 102.
- Ebe K, Bando H, Muneta T, Bando M, Yonei Y. Investigation of C-peptide Index for Carbo70 by the severity of diabetic state. *Clin Med Case Rep.* 2018; 2: 113.
- Hashimoto M, Ebe K, Bando H, Bando M, Muneta T. Response of insulin secretion to small amount of meal on low carbohydrate diet (LCD). *Biomed Sci J.* 2020; 1: 03. doi: [10.52588/biomedscij.1.03](https://doi.org/10.52588/biomedscij.1.03)
- Kanazawa S, Kato Y, Bando H, et al. Significant weight reduc-

tion by super low carbohydrate diet with improved lipid profiles. *International Medicine*. 2020; 2(3): 175-178. doi: [10.5455/im.116245](https://doi.org/10.5455/im.116245)

21. Ebe K, Hashimoto M, Bando H, Bando M, Muneta T. Proposal of Meal Tolerance Test (MTT) for investigating ability of insulin secretion for small carbohydrate load. *Diab Res Open Access*. 2020; 2(2): 31-37. doi: [10.36502/2020/droa.6166](https://doi.org/10.36502/2020/droa.6166)

22. Macedo RCO, Santos HO, Tinsley GM, Reischak-Oliveira A. Low-carbohydrate diets: Effects on metabolism and exercise - A comprehensive literature review. *Clin Nutr ESPEN*. 2020; 40: 17-26. doi: [10.1016/j.clnesp.2020.07.022](https://doi.org/10.1016/j.clnesp.2020.07.022)

23. Shankar SS, Vella A, Raymond RH, Staten MA, Calle RA, Bergman RN, et al. Standardized mixed-meal tolerance and arginine stimulation tests provide reproducible and complementary measures of  $\beta$ -cell function: Results from the foundation for the National Institutes of Health Biomarkers Consortium Investigative Series. *Diabetes Care*. 2016; 39: 1602-1613. doi: [10.2337/dc15-0931](https://doi.org/10.2337/dc15-0931)

24. Rijkkelijkhuizen JM, Girman CJ, Mari A, et al. Classical and model-based estimates of beta-cell function during a mixed meal vs. an OGTT in a population-based cohort. *Diabetes Res Clin Pract*. 2009; 83: 280-288. doi: [10.1016/j.diabres.2008.11.017](https://doi.org/10.1016/j.diabres.2008.11.017)

25. Paglialunga S, Guerrero A, Roessig JM, Rubin P, Dehn CA. Adding to the spectrum of insulin sensitive populations for mixed meal tolerance test glucose reliability assessment. *J Diabetes Metab Disord*. 2016; 15: 57. doi: [10.1186/s40200-016-0279-x](https://doi.org/10.1186/s40200-016-0279-x)

26. Park SY, Gautier JF, Chon S. Assessment of insulin secretion and insulin resistance in human. *Diabetes Metab J*. 2021; 45(5): 641-654. doi: [10.4093/dmj.2021.0220](https://doi.org/10.4093/dmj.2021.0220)

27. Yasuhara D, Naruo T, Nagai N, Tanaka M, Muranaga T, Nozoe S. Insulinogenic index at 15 min as a marker of nutritional rehabilitation in anorexia nervosa. *Am J Clin Nutr*. 2003; 77: 292-299. doi: [10.1093/ajcn/77.2.292](https://doi.org/10.1093/ajcn/77.2.292)

28. Esteghamati A, Ashraf H, Khalilzadeh O, et al. Optimal cut-off of homeostasis model assessment of insulin resistance (HOMA-IR) for the diagnosis of metabolic syndrome: third national surveillance of risk factors of non-communicable diseases in Iran (SuRFNCD-2007). *Nutr Metab (Lond)*. 2010; 7: 26. doi: [10.1186/1743-7075-7-26](https://doi.org/10.1186/1743-7075-7-26)

29. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model assessment: Insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia*. 1985; 28: 412-419. doi: [10.1007/BF00280883](https://doi.org/10.1007/BF00280883)

30. Choi CS, Kim MY, Han K, Lee MS. Assessment of  $\beta$ -cell function in human patients. *Islets*. 2012; 4: 79-83. doi: [10.4161/isl.19270](https://doi.org/10.4161/isl.19270)

31. Bando H, Iwatsuki N, Sakamoto K, Ogawa T. Satisfactory rapid response to xultophy associated with Meal Tolerance Test (MTT) by carbohydrate loading. *Asp Biomed Clin Case Rep*. 2021; 4(2): 145-152. doi: [10.36502/2021/ASJBCCR.6245](https://doi.org/10.36502/2021/ASJBCCR.6245)

32. Hijikata M, Higa M, Ichijo T, Hirose T. A comparison of meal tolerance test and oral glucose tolerance test for predicting insulin therapy in patients with gestational diabetes. *Food Nutr Res*. 2021; 65: 10.29219/fnr.v65.5490. doi: [10.29219/fnr.v65.5490](https://doi.org/10.29219/fnr.v65.5490)

33. Faber OK, Binder C. C-peptide response to glucagon. A test for the residual beta-cell function in diabetes mellitus. *Diabetes*. 1977; 26(7): 605-610. doi: [10.2337/diab.26.7.605](https://doi.org/10.2337/diab.26.7.605)