

Brief Research Report

Role of Low-Level Laser Therapy as an Adjuvant in Functional Rehabilitation of Patients with Joint Stiffness in Post-Burn Contracture

M. Vijayaragavan, MBBS¹; Ravi K. Chittoria, MS, MCh, DNB, MNAMS, PhD^{1*}; Neljo Thomas, MS¹; Nishad Kerakada, MS¹; Sanikommu D. Reddy, MBBS²; Mannoju Kumar S, MBBS²; Sri H. Reddy, MS¹

¹Department of Plastic Surgery, Jawaharlal Institute of Postgraduate Medical Education and Research (JIPMER), Pondicherry 605006, India

²Department of General Surgery, Jawaharlal Institute of Postgraduate Medical Education and Research (JIPMER), Pondicherry 605006, India

*Corresponding author

Ravi K. Chittoria, MS, MCh, DNB, MNAMS, PhD

Professor, Head of IT Wing and Telemedicine, Department of Plastic Surgery and Telemedicine, Jawaharlal Institute of Postgraduate Medical Education and Research (JIPMER), Pondicherry 605006, India; E-mail: drchittoria@yahoo.com

Article information

Received: December 20th, 2021; Revised: December 30th, 2021; Accepted: January 4th, 2022; Published: January 21st, 2022

Cite this article

Vijayaragavan M, Chittoria RK, Thomas N, et al. Role of low-level laser therapy as an adjuvant in functional rehabilitation of patients with joint stiffness in post-burn contracture. *Radiol Open J.* 2022; 6(1): 1-3. doi: [10.17140/ROJ-6-135](https://doi.org/10.17140/ROJ-6-135)

ABSTRACT

Introduction

Post-burn contracture is a known sequelae of burn injuries involving neck, axilla, elbow, and hand. There are many methods which are used as adjuvant in rehabilitation of post-burn contracture patients. The low-level laser therapy (LLLT) is one of the methods that can be used as an adjuvant in rehabilitation of joint stiffness following burns and but with very few data available from India. In this article we share our experience of using LLLT for functional rehabilitation of a patient with joint stiffness following post-burn contracture.

Materials and Methods

This study was conducted in a burns centre in a tertiary care institute in South India. Informed consent and departmental ethical committee clearance were obtained prior to study. The subject was a young boy with no known co-morbidities with stiffness of metacarpophalangeal and interphalangeal joint of index and middle finger of right hand. Patient was treated with contracture release with full thickness skin graft. The patient received low-level laser therapy to induce collagenolysis and to increase the movement of the joints. Each joint received laser therapy for duration of 125-second every time for 15-minutes for 2 sessions in a span of 4-days.

Result

LLLT has been found to be useful in early rehabilitation of post-burn contracture patients with joint stiffness. No side effects were observed during the study.

Conclusion

The LLLT is an effective, safe, and non-invasive adjuvant in functional rehabilitation of our patient with joint stiffness following post-burn contracture. We suggest further study including multiple subjects and a control group for validation of the result.

Keywords

Low-level laser therapy (LLLT); Post-burn contracture; Joint stiffness; Functional rehabilitation.

INTRODUCTION

Post-burn contracture is a known sequelae of burn injuries involving neck, axilla, elbow and hand. It occurs due to disorganized structure of the collagen in the wound areas. Various treatment modalities such as skin grafting are used to release the scars and to restore the function to the joint involved. Even though

these surgeries aid in anatomical restoration, functional restoration remains an important issue. After these reconstructive surgeries, patient may develop joint stiffness if post-operative physiotherapy is inadequate which can be due to various reasons such as pain, lost to follow-up, etc. There are many methods which are used as adjuvant in rehabilitation of post-burn contracture patients, one of them and with very few data available from India is low-level laser

therapy (LLLT). In this article we share our experience of using LLLT for functional rehabilitation of a patient with joint stiffness following post-burn contracture.

MATERIALS AND METHODS

This study was conducted in a burns centre under Department of Plastic Surgery in a Tertiary care institute in South India during the month of November- December 2021. Informed consent and Departmental Ethical Committee clearance were obtained prior to study. The subject was a young boy with no known co-morbidities with stiffness of metacarpophalangeal and interphalangeal joint of index and middle finger of right hand (Figure 1). Patient was treated with contracture release with full thickness skin graft (Figure 2). Patient developed stiffness of metacarpophalangeal joint and interphalangeal joint due to splinting following full thickness skin grafting (FTSG). The patient received LLLT to induce collagenolysis and to increase the movement of the joints.

ning mode (non-contact delivery) with 60 cm distance between laser source and the joint. Each joint received laser therapy for duration of 125-second every time for 15-minutes for 2 sessions in a span of 4-days (Figure 3).

Figure 1. Post-Burn Contracture Involving Multiple Fingers



Figure 2. Contracture Release Marking



Figure 3. Low-Level Laser Therapy Being Applied After PBC Release



RESULT

Low-level laser therapy has been found to be useful in early rehabilitation of post-burn contracture patients with joint stiffness. (Figures 4 and 5) No side effects were observed during the study.

Figure 4. Condition at Discharge with Near Normal Extension



Figure 5. Near Normal Flexion



Low-level laser source we used was Gallium Arsenide (gas) diode red laser of wavelength 650 nm, frequency 10 KHZ and output power 100 mw, which was a continuous beam laser with an energy density of 4 J/cm². Machine delivers laser in scan-

DISCUSSION

Burn injuries impart a considerable burden on health care system. The burns stand as a common cause of trauma related death second to road traffic accidents in both developing and developed country both acute and long-term events associated with burns patients compels patient to seek medical care. After resuscitation, which is aimed at survival of the victim, restoration to pre-injury status, and return to daily activities becomes important for victim and treating team.¹ A healed burn patient may be left with scars with varying degrees of functional issues and cause social stigma among victims.

Post-burn contracture develops because of healing by secondary intention may cause anatomical and functional morbidity to the patient. These patients may also develop mental illness due to the deformity. Rehabilitation in the form of surgery to restore the anatomy, functional rehabilitation in the form of physiotherapy and psychosocial counselling is necessary for the patient to allow him to lead a normal life.

Post-burn contracture leads to deformity due to faulty collagen. LLLT helps in preventing deformity by collagenolysis. This also helps in preventing joint stiffness following splinting.

The acronym “light amplification by stimulated emission of radiation (LASER)”, are defined by a power density at 1500 mW/cm^2.^{2,3} Energy used in LLLT is much less than the one used for cutting, and ablation therapy. LLLT is a form of phototherapy that employs electromagnetic radiation, that can generate enough energy for interacting with living tissues. It produces photochemical and photophysical effects without generation of heat, with consideration of re-establishing cell homeostasis. Essentially, light energy is delivered topically in controlled way which is absorbed by photo-absorbers (chromophores) that transform it into chemical energy.⁴

Positive effects include increased formation of granulation tissue and acceleration of tissue repair, wound contraction, inflammation, modulation, and pain reduction.⁴

As per literature, low-energy photoemissions given at a wavelength range of 600 nm to 900 nm accelerate cell proliferation and promote wound healing.⁵ Its action is thought to:

- Stimulate respiratory chain components promoting adenosine triphosphate (ATP) synthesis,⁶ and hence increase rate of mitoses and fibroblast numbers.⁶
- Stimulate collagen and elastin production.⁷
- Stimulate microcirculation with dilatation of the capillaries and neovascularisation.⁸
- Cause collagenolysis.⁹
- Liberate mediator of inflammation-histamine, serotonin and bradykinin and hence activate macrophages.

- Regenerate lymphatic vessels.

CONCLUSION

Low-level laser therapy is found to be useful in management of joint stiffness our patient following contracture release without any side effects. The limitation of the study it was done on a single subject. Hence the authors suggest that a study including multiple subjects with a control group and multiple centres with randomisation to validate the exact result.

CONFLICTS OF INTEREST

The authors declare that they have no conflicts of interest.

REFERENCES

1. Goel A, Shrivastava P. Post-burn scars and scar contractures. *Indian J Plast Surg.* 2010; 43(Suppl): S63-S71. doi: 10.4103/0970-0358.70724
2. Karu TI. Low-power laser therapy. In: Vo-Dinh T. *Biomedical Photonics Handbook*, ed. London, UK: CRC Press; 2003: 48. 7-20.
3. Baxter CD. *Therapeutic Lasers: Theory and Practice*. London, UK: Churchill Livingstone; 1994.
4. Andrade FSSD, Clark RMO, Ferreira ML. Effects of low-level laser therapy on wound healing. *Rev Col Bras Cir.* 2014; 41(2): 129-133. doi: 10.1590/s0100-69912014000200010
5. Lichtenstein D, Morag B. Low-level laser therapy in ambulatory patients with venous stasis ulcers. *Laser Therapy.* 1998; 11: 71-78.
6. Karu T. Molecular mechanism of therapeutic effect of low intensity laser irradiation. *Dokl Akad Nauk SSSR.* 1986; 291: 1245-1249.
7. Saperia D, Glassberg E, Lyons RF, et al. Demonstration of elevated type I and type III procollagen mRNA levels in cutaneous wounds treated with helium-neon laser. Proposed mechanism for enhanced wound healing. *Biochem Biophys Res Commun.* 1986; 138: 1123-1128. doi: 10.1016/s0006-291x(86)80399-0
8. Schindl A, Schindl M, Schindl L, et al. Increased dermal angiogenesis after low-intensity laser therapy for a chronic radiation ulcer determined by a video measuring system. *J Am Acad Dermatol.* 1999; 40: 481-484. doi: 10.1016/s0190-9622(99)70503-7
9. The effects of low-level laser therapy in extensor tendon injuries between zones 5-8. Web site. <https://clinicaltrials.gov/ct2/show/NCT04742296>. Accessed December 19, 2021.